

Occurrence and molecular characterization of *Giardia duodenalis* and *Cryptosporidium* spp. in sheep and goats reared under dairy husbandry systems in Greece

Nikolaos Tzanidakis, Smaragda Sotiraki, [...], and Thomas Geurden

Abstract

Giardia duodenalis and *Cryptosporidium* spp. are gastro-intestinal protozoa known to infect small ruminants. Both protozoa are also considered as a potential public health concern. The objective of this study was to determine their prevalence in lambs and goat kids kept under common Mediterranean dairy husbandry systems and to identify the species and genotypes infecting these small ruminants. In total, 684 faecal samples (429 from lambs and 255 from goat kids) were collected on 21 farms in Greece and examined using a quantitative immunofluorescence assay. *G. duodenalis* was detected in 37.3% of the lambs and 40.4% of the goat kids. On all but one of the farms *G. duodenalis* was detected. Most samples were typed as a mono-infection with *G. duodenalis* assemblage E, both on the β -giardin gene and the triose phosphate isomerase gene. Only 10% of samples were typed as mixed assemblage A and E infections. The prevalence of *Cryptosporidium* spp. was 5.1% in lambs and 7.1% in goat kids. In total, 8 out of the 14 farms with a sheep flock and 7 out of the 14 farms with a goat flock were positive. *Cryptosporidium parvum* (subtype IId), *C. ubiquitum* and *C. xiaoi* were identified, the latter especially in goat kids. In conclusion, the results of the present study illustrate that *G. duodenalis* and *Cryptosporidium* spp. occur frequently on both sheep and goats farms. The prevalence of zoonotic genotypes or species was low, indicating a limited but existing risk for zoonotic infections.

Keywords: *Giardia*, *Cryptosporidium*, Goat, Sheep, Dairy, Prevalence, Genotyping

Introduction

Giardia duodenalis and *Cryptosporidium* spp. are gastro-intestinal protozoa that affect a wide range of mammals. Both parasites have a direct life cycle and are known to cause enteritis. In small ruminants, mainly young lambs and goat kids are infected. The prevalence of both parasites in small ruminants varies considerably between studies worldwide [13]. The most common clinical symptoms associated with *G. duodenalis* are the excretion of malodorous, loose to diarrhoeic faeces and impaired weight gain, whereas *Cryptosporidium* spp. infection can lead to severe diarrhoea, depression, anorexia and weight loss [1, 12]. Mortality has been associated with cryptosporidiosis, especially in animals with concurrent infections.

Since the initial claims on its potential public health relevance, eight genotypes or so-called assemblages have been identified within *G. duodenalis* [25]. Assemblage A and B are considered to be zoonotic genotypes, affecting both humans and small ruminants [1, 12, 16, 23], whereas Assemblage E is considered to be specific to hoofed livestock and has been found to be the most prevalent assemblage in lambs and goat kids [8, 12]. The genus *Cryptosporidium* consists of 20 species and more than 40 genotypes. In sheep, *C. parvum*, *C. xiaoi* and *C. ubiquitum* are most frequently identified [9, 12, 27, 29, 35, 41] whereas mainly *C. parvum* and to a lesser extent *C. xiaoi* has been reported in goats [9, 12, 35]. Due to the potential clinical and zoonotic relevance, there is a need to better understand the presence and abundance of both parasites in sheep and goat flocks.

The aim of this study was to estimate the prevalence of *G. duodenalis* and *Cryptosporidium* spp. infection in sheep and goat dairy farms in Greece. The small ruminant sector is very important in the Mediterranean basin from an economic, social and ecological point of view [5], especially in Greece with approximately 15 million sheep and goats which are kept traditionally for milk under low-input systems [20]. The most commonly applied farming systems practised in Greece can be categorized as extensive farming [43], and there is limited information on prevalence and especially molecular characterization of both parasites in these systems. The prevalence rates reported range from 33.3% to 49.6% for *G. duodenalis* and from 4.4% to 55% for *Cryptosporidium* spp. in sheep and goats [21, 32].

Materials and methods

Study design

The study was designed as a cross-sectional study in a high sheep and goat density area in Greece (i.e. the island of Crete where more than 1.5 million animals are kept). The farms enrolled in this survey were selected according to the following criteria: (a) type of animal

on the farm (sheep, goats, or mixed sheep and goats), and (b) management practices applied (“intensive management system” where stocking rates are high and the young animals are reared indoors until weaning (30–40 days), and “extensive management system” where stocking rates are lower and young animals are reared with their mothers mostly on pasture). Each farm was visited on a single occasion in a 6-month period, and only animals between the age of 1 day and 10 weeks were considered for inclusion in the study. Sample size was calculated based on the number of expected births as an indicator of the number of animals on the farms: of the lambs, 5% of the expected births on each farm were sampled and of the goat kids, 10% of the expected births were sampled. Data on the type of water supply for the animals (public network supply, a private drill hole or natural well), the age of lambs or goat kids and the presence of diarrhoea were recorded.

Parasitological examination

The faecal samples were examined in the laboratory using a quantitative immunofluorescence assay (IFA; Merifluor *Cryptosporidium/Giardia* kit; Meridian Diagnostics Inc.), as follows: 1 g of faeces was suspended in tap water and sieved three times through a layer of surgical gauze to withhold large debris. Sedimentation for at least 30 min was followed by discarding the supernatant. The remaining sediment was centrifuged at 3000 rpm for 5 min. The sediment was re-suspended in 1 mL of tap water. After thorough vortexing, an aliquot of 20 µL was applied onto an IFA slide. The samples, including a negative and positive control sample, were left to dry completely. After staining and incubating slides in a dark humidified chamber (for 30 min at room temperature), the entire slide was examined at 400× magnification under a fluorescence microscope. A sample was considered positive if at least one, clearly recognizable *Cryptosporidium* oocyst or *Giardia* cyst was identified. The number of (oo)cysts per gram of faeces was obtained by multiplying the total number of (oo)cysts on the slide by 50.

Molecular characterization

Positive isolates for both parasites were selected for DNA extraction, using the QIAamp Stool Mini Kit (Qiagen), according to the manufacturer’s instructions, incorporating an extended initial step of five freeze-thaw cycles (freezing in liquid nitrogen for 5 min and heating at 95 °C for 5 min) in the protocol to maximize (oo)cyst lesion. The selection of the positive isolates aimed to include at least one positive sample per farm.

For the amplification of the *Cryptosporidium* spp., the 18S ribosomal DNA (18S rDNA) gene PCR protocol was used (previously described in [40]), as well as a PCR targeting the 70 kDa heat shock protein (HSP70, described in [28]). Subgenotyping of the *C. parvum* positive samples was performed using the 60 kDa glycoprotein (gp60) gene [34]. *G. duodenalis* positive samples were characterized using the β-giardin gene [24], and the triose phosphate isomerase (TPI) gene [11]. Amplification products were visualized on 1.5% agarose gels with ethidium bromide. A positive and negative (PCR water) control sample was included in each PCR. PCR products were purified using the Qiaquick purification kit (Qiagen) and fully sequenced using the BigDye Terminator V3.1 Cycle Sequencing Kit (Applied Biosystems). Sequencing reactions were analysed on a 3100 Genetic Analyzer (Applied Biosystems) and assembled with Seqman II (DNASTAR, Madison, WI, USA). Sequences were compared with known sequences by BLAST-analysis against the NCBI database.

Results

A total of 21 farms with either a sheep (n = 7) or goat (n = 7) flock, or a mixed sheep and goat flock (n = 7) were visited. On the majority of the farms (14/21) animals were reared under the intensive management system, whereas on 7 farms, goats and sheep were reared under the extensive management system (Table 1). In total, 684 faecal samples were examined, of which 429 samples were from lambs and 255 from goat kids. Mean lamb age was 5 weeks (range 4–8 weeks) and mean goat kid age was 6 weeks (range 4–9 weeks).

Table 1.
Overview of the 21 farms included in the study, with the number of sheep and/or goats kept on management system (Sampled = number of animals sampled on each farm).

Farm identification number	Management system	Sheep	
		Number of animals	Sampled
1	Intensive	203	17
7	Intensive	500	36
10	Intensive	400	26
11	Intensive	550	41
14	Intensive	370	13
16	Intensive	300	13
18	Intensive	750	34
21	Intensive	650	49

Table 1. Overview of the 21 farms included in the study, with the number of sheep and/or goats on the farms (N) and the type of management system (Sampled = number of animals sampled on each farm).

The prevalence of *G. duodenalis* was 37.3% (n = 160/429) in lambs and 40.4% (n = 103/255) in goat kids. In total, 13 out of the 14 farms (92.9%) with a sheep flock, and 14 out of the 14 farms (100%) with a goat flock, were positive for *G. duodenalis* with a minimum of 4 positive samples on each positive farm. The prevalence of *Cryptosporidium* spp. oocysts was 5.1% (n = 22/429) in lambs and 7.1%

($n = 18/255$) in goat kids. In total, 8 out of the 14 farms (57.1%) with a sheep flock, and 7 out of the 14 farms (50.0%) with a goat flock, were positive with a minimum of 1 positive sample on each positive farm.

Intensity of *G. duodenalis* cyst excretion ranged from 50 to 800,000 cysts per gram (cpg) of faeces in lambs with an average of 48,989 cpg, and from 50 to 900,000 cpg for goat kids with an average of 94,053 cpg. The excretion level for *Cryptosporidium* oocysts was low in lambs with an average of 9143 oocysts per gram of faeces (range 200–31,900 opg), yet was high in goat kids with an average excretion of 47,744 opg (range 200–551,000 opg).

A minimum of 1 positive sample was selected for DNA extraction per farm for *G. duodenalis* and *Cryptosporidium* spp., respectively. In total, 71 samples yielded a positive amplicon for *G. duodenalis* (Table 2). The majority of the samples were typed as a mono-infection with the ruminant-specific assemblage E, both on the β -giardin gene and the TPI gene. Only a limited number of samples were typed as mixed assemblage A and E infections, both in lambs and in goat kids. For *Cryptosporidium*, 24 samples yielded a positive amplicon (Table 3). Three different *Cryptosporidium* species (*C. parvum*, *C. ubiquitum* and *C. xiaoi*) were identified, although *C. xiaoi* was not identified in lambs. The *C. parvum* positive samples were typed as subtype IId on the gp60 gene (IIdA4G2T14 and IIdA4G3T13).

Table 2.

Results for molecular identification of *Giardia duodenalis* positive samples in goat kids and lambs, based on the B-giardin (= beta-giardin) or triose phosphate isomerase (TPI) gene (NA = no amplification; A = assemblage E and F).

Animal species	Number of samples	B-giardin	TPI
Goat kids	35	1 (2.9), 1 (2.9), 1 (2.9), 1 (2.9)	1 (2.9)
Lambs	41	1 (2.4), 1 (2.4), 1 (2.4), 1 (2.4)	1 (2.4)
Total	76	2 (2.6), 2 (2.6), 2 (2.6), 2 (2.6)	2 (2.6)

Table 2. Results for molecular identification of *Giardia duodenalis* positive samples in goat kids and lambs, based on the B-giardin (= beta-giardin) or triose phosphate isomerase (TPI) gene (NA = no amplification; A = assemblage ...

Table 3.

Results for molecular identification of *Cryptosporidium* positive samples in goat kids and lambs.

Animal species	Number of samples	<i>C. parvum</i>	<i>C. ubiquitum</i>	<i>C. xiaoi</i>
Goat kids	14	1 (7.1), 1 (7.1), 1 (7.1), 1 (7.1)	1 (7.1)	1 (7.1)
Lambs	10	1 (10.0), 1 (10.0), 1 (10.0), 1 (10.0)	1 (10.0)	1 (10.0)
Total	24	2 (8.3), 2 (8.3), 2 (8.3), 2 (8.3)	2 (8.3)	2 (8.3)

Table 3. Results for molecular identification of *Cryptosporidium* positive samples in goat kids and lambs.

Discussion

The present study illustrates that *G. duodenalis* is highly prevalent in both lambs and goat kids, as the parasite was detected in all but one of the sheep flocks and in all goat flocks. The high farm and animal prevalence is in line with previous studies in small ruminants in Europe [4, 13, 18]. The high *G. duodenalis* prevalence and the potential association with production losses [31, 44] require an appropriate level of awareness of this infection on those farms in terms of disease management and prevention of infection. In contrast to *G. duodenalis*, the prevalence of *Cryptosporidium* was lower than anticipated in both lambs and goat kids, probably due to the age range of the animals included in the present study. Nevertheless, the farm prevalence on the sheep (57.1%) and goat (50.0%) farms does suggest that *Cryptosporidium* is widespread and is a potential threat to the small ruminant population. In previous studies in Greece, *Cryptosporidium* has been associated with large outbreaks of diarrhoea in both sheep and goat flocks [14, 15, 37], similar to other major small ruminant rearing countries [6, 33].

Both for *G. duodenalis* and for *Cryptosporidium* spp. a potential public health threat has been suspected, based on the high prevalence of these parasites in small ruminants and on extrapolation of molecular insights from other animal species, such as cattle or companion animals. However, recent molecular data seem to suggest that small ruminants are mostly infected with non-zoonotic genotypes of *G. duodenalis* [12, 18, 38, 39, 46, 47] and *Cryptosporidium* spp. [38]. On the other hand, potentially zoonotic genotypes or species such as *G. duodenalis* assemblage A and B [1, 2, 12, 19, 26, 30, 41], *C. parvum* [3, 6, 12, 22, 46], *C. hominis* [17], *C. meleagridis* [42] and *C. ubiquitum* [10, 12, 45] have been reported in small ruminants. Furthermore, the identification of potentially zoonotic genotypes does not necessarily imply that transmission occurs. Recently, host-associated populations of *C. parvum* have been described using a multi-locus genotyping (MLG) approach, and *C. parvum* populations found in goats were even found to differ from bovine and sheep MLGs [7]. Whether this is a true host-specific phenomenon or just a matter of the level of isolation and opportunities for out-crossing is still to be discussed. Nevertheless, the contradicting molecular findings illustrate the difficulty of evaluating a potential public health threat based solely on genetic data without considering the epidemiological background and transmission of infection. Direct transmission of *Cryptosporidium* infection through bottle feeding or petting of animals on educational farms has been described before [38], but is probably an occasional route of infection. Although there is no direct evidence of transmission of *Cryptosporidium* and *G. duodenalis* infections from small ruminants to the human population via contaminated water, it is considered a threat. Furthermore, the detection of both parasites in outbreaks and in water screening is not routine practice in most countries, and large waterborne outbreaks might go

unnoticed. In a recent study in Spain, the prevalence of *Cryptosporidium* and *Giardia* in water was significantly higher in the inland area, with higher concentration of livestock and fewer water treatment plants [4], illustrating that a variety of factors define the odds for infection. In the specific study area on the island of Crete, only a limited number of water basins are used over the island to produce drinking water for the local population and for the tourist population in the summer. The pastures surrounding the drinking water basins are all grazed by small ruminants. Whether these conditions lead to a substantial public health threat will need to be evaluated further in a longitudinal study, including sampling of water.

In goats, a large proportion of the *Cryptosporidium* positive samples were typed as *C. xiaoi*, both on 18S and HSP-70. This is in agreement with previous studies in Spain [6] and France [36], yet contradicts the initial claim that *C. xiaoi* infections are largely restricted to sheep [9]. In the current study, *C. xiaoi* was found in goats from three different farms, of which 2 farms maintained a goat-only flock and 1 farm managed a mixed flock. This illustrates that, although the introduction in the goat flocks might be due to contact with infected sheep, a *C. xiaoi* infection is easily maintained in goats. As advocated by Fayer and Santin [9], further epidemiological data will be needed to confirm whether the reports of *C. xiaoi* in goats are incidental or a regularly observed finding.

In conclusion, a high animal and farm prevalence of *G. duodenalis*, and a high farm prevalence of *Cryptosporidium* spp. were detected in both lambs and goat kids. Although mainly non-zoonotic species were identified in the present study, the frequent contact and proximity of grazing grounds to the natural water sources used to produce drinking water in the study area warrant further investigation of the public health relevance of these infections.

Acknowledgments

The authors would like to thank the farmers and research staff who participated in this study, with special thanks to Dr. Alexandros Stefanakis. The study was conducted within the framework of the COST Action FA0805 CAPARA. “Goat-parasite interactions: from knowledge to control”.

Notes

Cite this article as: Tzanidakis N, Sotiraki S, Claerebout E, Ehsan A, Voutzourakis N, Kostopoulou D, Stijn C, Vercruysse J & Geurden T: Occurrence and molecular characterization of *Giardia duodenalis* and *Cryptosporidium* spp. in sheep and goats reared under dairy husbandry systems in Greece. *Parasite*, 2014, **21**, 45.

Footnotes

★ **Novel Approaches to the Control of Parasites in Goats and Sheep.**

Invited editors: Hervé Hoste, Smaragda Sotiraki, and Michel Alvinerie

Article information

Parasite. 2014; **21**: 45.

Published online Sep 5, 2014. doi: [10.1051/parasite/2014048](https://doi.org/10.1051/parasite/2014048)

PMCID: PMC4154256

Nikolaos Tzanidakis,¹ Smaragda Sotiraki,¹ Edwin Claerebout,² Amimul Ehsan,² Nikolaos Voutzourakis,¹ Despoina Kostopoulou,¹ Casaert Stijn,² Jozef Vercruysse,² and Thomas Geurden^{2,*}

¹ Veterinary Research Institute – Hellenic Agricultural Organization Demeter 57001 Thessaloniki Greece

² Laboratory for Parasitology, Faculty of Veterinary Medicine, Ghent University Salisburylaan 133 9820 Merelbeke Belgium

*Corresponding author: Email: thomasgeurden@yahoo.com

Received December 29, 2013; Accepted August 25, 2014.

Copyright © N. Tzanidakis et al., published by EDP Sciences, 2014

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Articles from *Parasite* are provided here courtesy of **EDP Sciences**

References

1. Aloisio F, Filippini G, Antenucci P, Lepri E, Pezzotti G, Caccio S-M, Pozio E. 2006. Severe weight loss in lambs infected with *Giardia duodenalis* assemblage B. *Veterinary Parasitology*, 142, 154–158 [[PubMed](#)]
2. Berrilli F, D'Alfonso R, Giangaspero A, Marangi M, Brandonisio O, Kaboré Y, Glé C, Cianfanelli C, Lauro R, Di Cave D. 2012. *Giardia duodenalis* genotypes and *Cryptosporidium* species in humans and domestic animals in Côte d'Ivoire: occurrence and evidence for environmental contamination. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 106, 191–195 [[PubMed](#)]
3. Cacciò SM, Sannella AR, Mariano V, Valentini S, Berti F, Tosini F, Pozio E. 2013. A rare *Cryptosporidium parvum* genotype associated with infection of lambs and zoonotic transmission in Italy. *Veterinary Parasitology*, 191, 128–131 [[PubMed](#)]
4. Castro-Hermida JA, García-Preseido I, Almeida A, González-Warleta M, Correia Da Costa JM, Mezo M. 2011. *Cryptosporidium* spp. and *Giardia duodenalis* in two areas of Galicia (NW Spain). *Science of the Total Environment*, 409, 2451–2459 [[PubMed](#)]
5. De Rancourt M, Fois N, Lavin MP, Tchakerian E, Vallerand F. 2006. Mediterranean sheep and goat production: an uncertain future. *Small Ruminant Research*, 62, 167–179
6. Díaz P, Quilez J, Robinson G, Chalmers RM, Díez-Baños P, Morondo P. 2010. Identification of *Cryptosporidium xiaoi* in diarrhoeic goat kids (*Capra hircus*) in Spain. *Veterinary Parasitology*, 172, 132–134 [[PubMed](#)]
7. Drumo R, Widmer G, Morrison LJ, Tait A, Grelloni V, D'Avino N, Pozio E, Cacciò SM. 2012. Evidence of host-associated populations of *Cryptosporidium parvum* in Italy. *Applied Environmental Microbiology*, 78, 3523–3529 [[PMC free article](#)] [[PubMed](#)]
8. Ey PL, Mansouri M, Kulda J, Nohynkova E, Monis PT, Andrews RH, Mayrhofer G. 1997. Genetic analysis of *Giardia* from hoofed farm animals reveals artiodactyls-specific and potentially zoonotic genotypes. *Journal of Eukaryotic Microbiology*, 44, 625–635 [[PubMed](#)]
9. Fayer R, Santin M. 2009. *Cryptosporidium xiaoi* n. sp. (Apicomplexa: *Cryptosporidiidae*) in sheep (*Ovis aries*). *Veterinary Parasitology*, 164, 192–200 [[PubMed](#)]
10. Fiuza VR, Cosendey RI, Frazão-Teixeira E, Santin M, Fayer R, de Oliveira FC. 2011. Molecular characterization of *Cryptosporidium* in Brazilian sheep. *Veterinary Parasitology*, 175, 360–362 [[PubMed](#)]
11. Geurden T, Geldhof P, Levecke B, Martens C, Berkvens D, Casaert S, Vercruysse J, Claerebout E. 2008. Mixed *Giardia duodenalis* assemblage A and E infections in calves. *International Journal for Parasitology*, 38, 259–264 [[PubMed](#)]
12. Geurden T, Thomas P, Casaert S, Vercruysse J, Claerebout E. 2008. Prevalence and molecular characterization of *Cryptosporidium* and *Giardia* in lambs and goat kids in Belgium. *Veterinary Parasitology*, 155, 142–145 [[PubMed](#)]
13. Geurden T, Vercruysse J, Claerebout E. 2010. Is *Giardia* a significant pathogen in production animals? *Experimental Parasitology*, 124, 98–106 [[PubMed](#)]
14. Giadinis ND, Panousis NK, Papadopoulos E, Antoniadou-Sotiriadou K, Karatzias H. 2006. The effects of halofuginone in cases of cryptosporidiosis in newborn lambs and kids in Greece. *Proceedings of the 10th Greek Veterinary Congress, Athens*: p. 195–196
15. Giadinis N, Papadopoulos E, Panousis N, Papazahariadou M, Lafi SQ, Karatzias H. 2007. Effect of halofuginone lactate on treatment and prevention of lamb cryptosporidiosis: an extensive field trial. *Journal of Veterinary Pharmacology and Therapeutics*, 30, 578–582 [[PubMed](#)]
16. Giangaspero A, Paoletti B, Iorio R, Traversa D. 2005. Prevalence and molecular characterization of *Giardia duodenalis* from sheep in central Italy. *Parasitology Research*, 96, 32–37 [[PubMed](#)]
17. Giles M, Chalmers R, Pritchard G, Elwin K, Mueller-Doblies D, Clifton-Hadley F. 2009. *Cryptosporidium hominis* in a goat and a sheep in the UK. *Veterinary Record*, 164, 24–25 [[PubMed](#)]
18. Gómez-Muñoz MT, Navarro C, Garijo-Toledo MM, Dea-Ayuela MA, Fernández-Barredo S, Pérez-Gracia MT, Domínguez-Márquez MV, Borrás R. 2009. Occurrence and genotypes of *Giardia* isolated from lambs in Spain. *Parasitology International*, 58, 297–299 [[PubMed](#)]
19. Gómez-Muñoz MT, Cámara-Badenes C, Martínez-Herrero Mdel C, Dea-Ayuela MA, Pérez-Gracia MT, Fernández-Barredo S, Santin M, Fayer R. 2012. Multilocus genotyping of *Giardia duodenalis* in lambs from Spain reveals a high heterogeneity. *Research in Veterinary Science*, 93, 836–842 [[PubMed](#)]
20. Hadjigeorgiou I, Vallerand F, Tsimpoukas K, Zervas G. 2002. The socio-economics of sheep and goat farming in Greece and the implications for the future rural development. *Options Méditerranéennes Series B*, 39, 83–93

21. Himonas CA, Antoniadou-Sotiriadou KS, Sotiraki ST, Papazahariadou MG. 1998. Intestinal protozoa of animals in Macedonia. Bulletin of the Hellenic Veterinary Medical Society, 49, 300–306
22. Imre K, Luca C, Costache M, Sala C, Morar A, Morariu S, Ilie MS, Imre M, Dărăbuș G. 2013. Zoonotic *Cryptosporidium parvum* in Romanian newborn lambs (*Ovis aries*). Veterinary Parasitology, 191, 119–122 [PubMed]
23. Karanis P, Ey PL. 1998. Characterization of axenic isolates of *Giardia intestinalis* established from humans and animals in Germany. Parasitology Research, 84, 442–449 [PubMed]
24. Lalle M, Jimenez-Cardosa E, Cacciò SM, Pozio E. 2005. Genotyping of *Giardia duodenalis* from humans and dogs from Mexico using a beta-giardin nested polymerase chain reaction assay. Journal of Parasitology, 91, 203–205 [PubMed]
25. Lasek-Nesselquist E, Welch DM, Sogin ML. 2010. The identification of a new *Giardia duodenalis* assemblage in marine vertebrates and a preliminary analysis of *G. duodenalis* population biology in marine systems. International Journal for Parasitology, 40, 1063–1074 [PMC free article] [PubMed]
26. Lim YA, Mahdy MA, Tan TK, Goh XT, Jex AR, Nolan MJ, Sharma RS, Gasser RB. 2013. First molecular characterization of *Giardia duodenalis* from goats in Malaysia. Molecular and Cellular Probes, 27, 28–31 [PubMed]
27. McLauchlin J, Amar C, Pedraza-Diaz S, Nichols GL. 2000. Molecular epidemiological analysis of *Cryptosporidium* spp. in the United Kingdom: Results of genotyping *Cryptosporidium* spp. in 1,705 faecal samples from humans and 105 faecal samples from livestock animals. Journal of Clinical Microbiology, 38, 3984–3990 [PMC free article] [PubMed]
28. Morgan UM, Monis PT, Xiao L, Limor J, Sulaiman I, Raidal S, O'Donoghue P, Gasser R, Murray A, Fayer R, Blagburn BL, Lal AA, Thompson RC. 2001. Molecular and phylogenetic characterisation of *Cryptosporidium* from birds. International Journal for Parasitology, 31, 289–296 [PubMed]
29. Mueller-Doblies D, Giles M, Elwin K, Smith RP, Clifton-Hadley FC, Chalmers RM. 2008. Distribution of *Cryptosporidium* species in sheep in the UK. Veterinary Parasitology, 154, 214–219 [PubMed]
30. Nolan MJ, Jex AR, Pangasa A, Young ND, Campbell AJ, Stevens M, Gasser RB. 2010. Analysis of nucleotide variation within the triose-phosphate isomerase gene of *Giardia duodenalis* from sheep and its zoonotic implications. Electrophoresis, 31, 287–298 [PubMed]
31. Olson ME, McAllister TA, Deselliers L, Morck DW, Cheng KJ, Buret AG, Ceri H. 1995. Effects of giardiasis on production in a domestic ruminant (lamb) model. American Journal of Veterinary Research, 56, 1470–1474 [PubMed]
32. Panousis N, Diakou A, Giadinis N, Papadopoulos E, Karatzias H, Haralampidis S. 2008. Prevalence of *Cryptosporidium* infection in sheep flocks with a history of lambs' diarrhoea. Revue de Médecine Vétérinaire, 159, 528–531
33. Paraud C, Pors I, Chartier C. 2010. Evaluation of oral tilmosin efficacy against severe cryptosporidiosis in neonatal kids under field conditions. Veterinary Parasitology, 170, 149–152 [PubMed]
34. Peng MM, Matos O, Gatei W, Das P, Stantic-Pavlinic M, Bern C, Sulaiman IM, Glaberman S, Lal AA, Xiao L. 2001. A comparison of *Cryptosporidium* subgenotypes from several geographic regions. Journal of Eukaryotic Microbiology Supplement, 48, 28S–31S [PubMed]
35. Quilez J, Torres E, Chalmers RM, Hadfield SJ, Cacho E, Sanchez-Acedo C. 2008. *Cryptosporidium* Genotypes and subtypes in lambs and goat kids in Spain. Applied and Environmental Microbiology, 74, 6026–6031 [PMC free article] [PubMed]
36. Rieux A, Paraud C, Pors I, Chartier C. 2013. Molecular characterization of *Cryptosporidium* spp. in pre-weaned kids in a dairy goat farm in western France. Veterinary Parasitology, 192, 268–272 [PubMed]
37. Robertson LJ. 2009. *Giardia* and *Cryptosporidium* infections in sheep and goats: a review of the potential for transmission to humans via environmental contamination. Epidemiology and Infection, 137, 913–921 [PubMed]
38. Robertson LJ, Gjerde BK, Hansen EF. 2010. The zoonotic potential of *Giardia* and *Cryptosporidium* in Norwegian sheep: a longitudinal investigation of 6 flocks of lambs. Veterinary Parasitology, 171, 140–145 [PubMed]
39. Ruiz A, Foronda P, Gonzalez JF, Guedes A, Abreu-Acosta N, Molina JM, Valladares B. 2008. Occurrence and genotype characterization of *Giardia duodenalis* in goat kids from the Canary Islands, Spain. Veterinary Parasitology, 154, 137–141 [PubMed]
40. Ryan UM, Bath C, Rovertson I, Read C, Elliot A, Mcinnes L, Traub R, Besier B. 2005. Sheep may not be an important zoonotic reservoir for *Cryptosporidium* and *Giardia* Parasites. Applied Environmental Microbiology, 71, 4992–4997 [PMC free article] [PubMed]
41. Santin M, Trout JM, Fayer R. 2007. Prevalence and molecular characterization of *Cryptosporidium* and *Giardia* species and genotypes in sheep in

Maryland. *Veterinary Parasitology*, 146, 17–24 [[PubMed](#)]

42. Silverlås C, Mattsson JG, Insulander M, Lebbad M. 2012. Zoonotic transmission of *Cryptosporidium meleagridis* on an organic Swedish farm. *International Journal for Parasitology*, 42, 963–967 [[PubMed](#)]

43. Stefanakis A, Volanis M, Zoiopoulos P, Hadjigeorgiou I. 2007. Assessing the potential benefits of technical intervention in evolving the semi-intensive dairy-sheep farms in Crete. *Small Ruminant Research*, 72, 66–72

44. Sweeny JP, Robertson ID, Ryan UM, Jacobson C, Woodgate RG. 2011. Impacts of naturally acquired protozoa and strongylid nematode infections on growth and faecal attributes in lambs. *Veterinary Parasitology*, 184, 298–308 [[PubMed](#)]

45. Wang Y, Feng Y, Cui B, Jian F, Ning C, Wang R, Zhang L, Xiao L. 2010. Cervine genotype is the major *Cryptosporidium* genotype in sheep in China. *Parasitology Research*, 106, 341–347 [[PubMed](#)]

46. Yang R, Jacobson C, Gordon C, Ryan U. 2009. Prevalence and molecular characterisation of *Cryptosporidium* and *Giardia* species in pre-weaned sheep in Australia. *Veterinary Parasitology*, 161, 19–24 [[PubMed](#)]

47. Zhang W, Zhang X, Wang R, Liu A, Shen Y, Ling H, Cao J, Yang F, Zhang X, Zhang L. 2012. Genetic characterizations of *Giardia duodenalis* in sheep and goats in Heilongjiang Province, China and possibility of zoonotic transmission. *PLoS Neglected Tropical Diseases*, 6, e1826. [[PMC free article](#)] [[PubMed](#)]